# COURI CENTER

## T by Couri

## Post Insertion Instructions for Men

- A. Your insertion site has been closed with a dissolvable suture and covered with a bandage.
  - 1. The outer layer is the pressure bandage consisting of folded gauze and wide paper tape. Please remove the bandage in 48 hrs.
  - 2. You can remove the suture after 7-10 days. If you would prefer the suture be removed here, please call the office.
- B. THREE Day Rule:
  - 1. Please avoid the following for three (3) days:
    - Tub Baths
- Hot Tubs or Swimming
  Stronuous lifting
- Soaking in water

- Vigorous exercise Strenuous lifting
- 2. Showers are fine. Moderate exercise after 24 hrs is fine.

#### C. Other Important Notes:

- 1. Avoid scrubbing the site until the incision is well healed (about 7 days)
- 2. The insertion site may be uncomfortable for up to 2 to 3 weeks.
- 3. You may experience bruising, swelling, and/or redness of the insertion site which may last from a few days up to 2 to 3 weeks.
- 4. You may notice some pinkish or bloody discoloration on the outer bandage. This is normal.
- 5. If you experience bleeding from the incision, apply firm pressure for 5 minutes.
- Ice your incision area three (3) times a day for ten (10) minutes at a time on the day of your pellet insertion.
  This will help reduce any swelling, bruising or discomfort that you may experience.
- 7. You may take Tylenol or Ibuprofen as needed if you experience any discomfort.
- D. Please call us if you have any active bleeding or pus coming from the incision site.

#### E. Reminders

- Please have your labs rechecked in:
  □ 2-3 weeks before your next pellet insertion for accuracy of dosing.
- Other office requests:
  □ Yearly Prostate Exam
- 3. Please stay up to date with your primary care physician and yearly physicals.